actical Combat Casualty laye

August 2010



Direct from the Battlefield: TCCC Lessons Learned in Iraq and Afghanistan

Learned in Iraq and

- Afghanistan
 Reports from Joint Theater Trauma
 System (JTTS) weekly Trauma Telecons
 - Every Thursday morning worldwide telecon to discuss every serious casualty from that week
- Published medical re
- · Armed Forces Medica Examiner's Office re
- Feedback from docto corpsmen, medics, and PJs





Train ALL Combatants in TCCC

- Potentially preventable deaths averaging about 20% of all fatalities
- Units that train all members in TCCC have drastically reduced this incidence
- Need to train <u>ALL</u> combatants in TCCC



Hemorrhage

This casualty was wounded by an RPG explosion and sustained a traumatic amputation of the right forearm at the midforearm level and a right leg wound. He bled to death from his leg wound despite the placement of three fold-expedient

tourniquets.

What could have saved him

C.A.T. Tourniquet TCCC training for

<u>all</u>

unit members

*Note: Medic killed

<u>at</u>



- Get tourniquets on BEFORE onset of shock
 - Mortality is very high if casualties already in shock before tourniquet application
- If bleeding is not controlled and distal pulse not eliminated with first tourniquet use a second one installed.

proximal to first

 Increasing the tourniquet V with a second tourniquet controls

ghanistan - Nov 2009

- Soldier with gunshot wound to left leg
- Open fracture left femur
- Injury to popliteal artery and vein
- Three CAT tournique placed
- Life saved
- Leg doing well
- 2-3 casualties/week saved with tournic



 Tighten velcro band on tourniquets as tight as possible before starting to use windlass - a loose velcro band contributes to tourniquet malfunction

· Should be effective with approximately three degree turns of win

Use second tournique needed

To urniquets

 Fake CAT tourniquets that are prone to malfunction are turning up in theater - ensure that you have this NSN tourniquet:

· NSN 6515 01 521 7076









DEFENSE LOGISTICS AGENCY

DEFENSE SUPPLY CENTER PHILADELPHIA 700 ROBBINS AVENUE PHILADELPHIA, PENNSYLVANIA 19111-5092

DSCP-FSFB 10-150 April 14, 2010

MEMORANDUM FOR USAMMA, NAVMEDLOGCOM, AFMLO, MARCORSYSCOM, DMMPO.

SUBJECT: QUALITY ASSURANCE URGENT PRODUCT SAFETY ALERT.

1. REFERENCES:

A. ITEM: Tourniquet, Nonpneumatic; C-A-Tourniquet®. NSN 6515-01-521-7976.

B. Item No(s): NAR-CAT, 30-0001 Serial/Lot No(s): N/A

C. Manufacturer: Composite Resources, Inc., 485 Lakeshore Parkway, Rock Hill, SC

D. Distributors:

North American Rescue Inc., 35 Tedwall Court, Green, SC;

Cardinal Health, 1430 Waukegan Road, McGaw Park, IL.

Owens and Minor, 9120 Lockwood Blvd, Mechanicsville, VA:

American Purchasing Services (DBA American Medical Depot) 4380 NW 135th St, Opa Locka, FL;

Phoenix Textile Corporation, 21 Commerce Drive, O'Fallon, MO...

E. Authorized for procurement through DoD Supply Chain Only.

SAFETY ALERT: CRITICAL LIFE-SAVING ITEM.



Counterfeit CAT Tourniquets

SAFETY ALERT: CRITICAL LIFE-SAVING ITEM.

A. REASON: DLA has become aware of similar products manufactured to closely resemble the C-A-Tourniquet® and available for purchase through non-DoD websites. Authorized DoD procurement gateways will supply only the approved commercial part from authorized distributors. These products were first encountered several years ago in a depot in Afghanistan and thought to have been purged from the system. They were then of obvious inferior construction and quite recognizable as a substitute for the real thing. Today the product is very difficult to distinguish from the C-A-Tourniquet® down to duplicate markings and symbols.

Although there is no direct evidence against these duplicate products, several reports indicate that they are of inferior design and may cause serious injury or death.

B. RECOMMENDED STRATEGY: The above distributors, supplying the Composite Resources product exclusively, are the only authorized source for this device.

The FDA regulates this product as a Class 1 device, which means that there is no requirement for a premarket notification application and FDA clearance is not required before marketing the device in the U.S. However, these manufacturers are required to register their establishment with FDA.

If you have purchased these devices from any other source, it is recommended that they be suspended from use and replaced by the recommended product. Please report suspended quantities to your logistical supply office.

Some examples of non-authorized Internet sources for duplicate product that may be hazardous are: www.world-element.com; ID No. EX 159; and


http://www.airsoftglobal.com/product_info.php?products_id=11454; JD EL-ACC-EX159-AG.

CAT (GEN III) vs. F-CAT

Executive Summary

Introduction:

- The Element Cat (E-CAT) is a very carefully made counterfeit CAT tourniquet.
- It is manufactured in Hong Kong for \$8.50 (USD) per item.
- 3. There are no limits to the number that can be purchased.
- They are available on the internet, and anyone can purchase them.
- They were designed to look, feel and act like a CAT (GEN III).
- They ARE a counterfeit tourniquet.





CAT (GEN III) vs. F-CAT

Executive Summary

CAT (GEN III)

Package from NARP, Inc. Looks nothing like the F-CAT package.

E-CAT:

Packaged in plastic bag with paper top.

The sticker on the bag call the tourniquet the "Combat Application Tourniquet" and lists the NSN assigned to NARP.



E-CAT





CAT Generation VI

10/06/2009

Length of Tourniquet changed to 37 ½" Manufacturer and Lot Stamp with date manufactured added to the strap





2009 Officer Kim

Munley

- 12 dead; 31 wounded on 5 Nov 09
- Officer Munley got shooter; shot in both thighs
- Direct pressure and makeshift tourniquets used by several physicians unsuccessful at controlling hemorrhage
 - went into shock
- Saved by Army 68W medic with a CAT tourniquet on left thigh



Tourniquet on Uninjured Arm

- · JTTS Trauma Telecon 8 April 2010
- IED casualty
- Arrived at Kandahar with CAT in place on left arm
- · Evaluation: no injuries sustained on left arm
- · Follow-up: No explanation available
- · Lessons Learned:
 - · No injury No tourniquet
 - · Remember to reassess your casualties



Wear Your Eye Protection!

- · Jan 2010
- 22 y/o near IED without eye protection
- Now blind in both eyes





Without eye pro - both eyes being re

th eye pro - eyes OK



Penetrating Eye Trauma

- Rigid eye shield for obvious <u>or suspected</u> eye wounds - often not being done -SHIELD AND SHIP!
- Not doing this may cause permanent loss of vision - use a shield for <u>any</u> injury in or around the eye

always

Shield after injury



No shield after injury



Eye Protection





- Use your tactical eyewear to cover the injured eye if you don't have a shield.
- Using tactical eyewear in the field will

JTTS Trauma Telecon 9 Sept 2010

- · Recent case of endophthalmitis
- Reminder shield and moxifloxacin in the field for penetrating eye injuries
- · Also need to continue moxi both topically an systemically in the M7
- Many antibiotics <u>do no</u> <u>penetrate well</u> into the eye

Patched Open Globe 22 July 2010

- Shrapnel in right eye from IED
- · Had rigid eye shield placed
- Reported as both pressure patched and as having a gauze pad placed under the fox shield without pressure
- Extruded uveal tissue noted at time of operative repair of globe
- No gauze! COL Robb Mazzoli: Gauze can adhere to iris tissue and cause further extrusion when removed <u>even if no pressure is applied to eye.</u>



Surgical Airways

Joint Theater Trauma System Email 24 September 09

- 3 field crics done incorrectly in OIF
- · One through center of thyroid cartilage and through one of the vocal co

Thyroid cartilage
Incision site
Cricoid cartilage



Surgical Airways: The Rest of the Story

"The setting of the casualty care was at night in a nonpermissive environment. The medic had sustained a sacral injury and damaged his NVG's during a hard landing on infil. The casualty had sustained a gunshot wound to the jaw. The medic was not called to the scene for ten minutes due to an ongoing firefight. The jaw was shattered and he had heavy maxillofacial bleeding. The recovery position was attempted repeatedly, but the casualty refused to remain like that. Anxiolysis was attempted with Versed to facilitate maintaining the airway with position alone, but did not work. The casualty became increasingly combative and the decision was made to perform the cric out of fear of completely losing the airway during evacuation. Due to the fact that the medic's NVGs were damaged, an operator (former 18D with two successful prior combat cric's) attempted the precedure with assistance by the



Surgical Airways

Recommendations:

- Live tissue training for this procedure if possible
- "Sim Man" trainer may be secondbest option
- Don't attempt surgical airway just because the casualty is uncon
- Try the "sit-up and lean forwa position prior to attempting a airway



Surgical Airways

If you cut the endotracheal Tube, you must tape it very securely or the tube will slip down into the trachea, cease to function correctly, and have surgically remove

Like this one.....



IED Casualties

- IED blast casualties often have multiple mechanisms of injury
 - Blunt trauma
 - Penetrating trauma
 - · Blast
 - · Burns
- Majority of casual are now from Il





IED Casualties

· IED casualties - many have spinal fractures,

especially thoracic

Try to maintain spinal alignment in

blunt

trauma casualti





 IED events - be alert for secondary IEDs or ground assaults after initiation of the IED



Aviation Personnel Need CCC?

In-Flight Tourniquet 24 June 2010

- AF Pave Hawk pilot on EVAV mission to pick up wounded UK soldier
- GSW both legs
- Severe bleeding R leg
- PJ crawled up into cockpit and applied tourniquet
- Bleeding controlled pilot completed mission

JTTS Trauma Telecon 26 Aug 2010

- 23 y/o male
- · GSW left infraclavicular area with external hemorrhage
- "Progressive deterioration"
- External hemorrhage noted to increase as casualty resuscitated in ED
- No record of Combat Gauze use
- · All injuries noted to be extrapleural
- · Lesson learned: see following slide





It doesn't work if you don't

FEEDBACK TO THE FIELD:

Perforation of the Sternum by an Intraosseous Infusion Device


H T Harcke, COL, MC, USA Chief, Forensic Radiology Armed Forces Institute of Pathology

E Mazuchowaski, Lt Col (Sel), USAF, MC Deputy Medical Examiner Office of the Armed Forces Medical Examiner

CASE OVERVIEW

- IED detonated in the decedent's vicinity.
- Catastrophic injury to the lower extremities and pelvis, to include traumatic amputation of the lower legs.
- Emergency treatment included tourniquets, sternal IO-IV, and proximal humeral IO-IV's.






Note sternal IO in place

Autopsy CT Scan

Sagittal MDCT image shows the IO-IV needle passes through the sternum with the tip in the anterior mediastinum (arrow).



s is NOT where you want the infused fluids to

Comparison of the devices:

Note size, color and packaging differences.







Do you really want to try to tell these two IO needles apart in the dark in a

Comparison of the devices:

Note size, color and packaging differences.







Do you really want to try to tell these two IO needles apart in the dark in a





Questions?